# Impact of Psychosocial Health on Pregnancy Intention

Pamela J. Maxson, Ph.D.

Christina Gibson-Davis, Ph.D., Geeta Swamy, M.D., Redford Williams, M.D., Marie Lynn Miranda, Ph.D.






# Children's Environmental Health Initiative





# Relationships among SCEDDBO Projects





# **Study Design**

- Prospective cohort study of pregnant women in Durham, NC
- Psychosocial assessment through surveys
- Maternal blood draws at 28 weeks (genetics) and at birth (environmental exposures)
- Delivery samples: placental tissue, cord segment, cord blood
- Pregnancy outcomes including:
  - Birth weight
  - Gestation
  - Maternal medical complications




### **Study Participants**

- Over 18 years of age
- Speak/read English





- Singleton gestation, no known congenital abnormalities
- Between 18-28 weeks gestation at recruitment
- Recruitment (current=748; target=1500)
  - Duke OB clinic (409, 121 NHW, 249 NHB, 26 H)
  - Lincoln Community Health Center-Durham County Health Department (339, 20 NHW, 272 NHB, 33 H)






# **Demographics of Participants**









### **Pregnancy Intention Survey Question**

- Think back to just before you were pregnant with this child. How did you feel about becoming pregnant?
  - I wanted to be pregnant at that time.
  - I wanted to be pregnant sooner.
  - I wanted to be pregnant later. Later
  - I didn't want to be pregnant at that time or at any time in the future.

Never

Now



# **Pregnancy Intention Status**

- Now
  - N=240
  - 39% of participants
- Later
  - N=260
  - 42% of participants
- Never
  - N=113
  - 19% of participants

61% of the pregnancies are mistimed or unwanted



# **Psychosocial Instruments**

- CES-D
- John Henryism
- ISEL
- Paternal Support
- Self-Efficacy
- Perceived Stress
- NEO





# Depression (CES-D)

Lowest depressive symptoms

Highest depressive symptoms

Now Later Never

$$p = < .08$$

p<.0001



# **ISEL- Interpersonal Support**

# Most support Least support

Now Later Never


p<.08

p<.07

p<.01



# **Paternal Support**





# Self-efficacy

Highest Self-efficacy

Now

Later

Never

p<.05

p<.05



#### **Perceived Stress**

# Now Later Never p<.0001 p<.0001



# **Extraversion- NEO**

#### Most extraverted

Least extraverted

Now Later Never

p<.08

p<.07

p<.01



# **Neuroticism- NEO**

Least neuroticMost neuroticNowLaterNever

p<.001

p<.001

p<.001



# **Personality- NEO**

- -Openness to Experience
- -Agreeableness
- -Conscientiousness

Highest Lowest

Now Later Never

**Significant** 

Significant



#### **Maternal Behaviors**

- No differences in tobacco usage between intention statuses.
- No differences in alcohol usage between intention statuses.



# **Pregnancy Outcomes**

- Birth weight
- Gestational age at delivery
- Prevalence of gestational diabetes

Pregnancy related hypertension

	Gestational
Intention	Age
Never*	37.6 weeks
Later	38.2 weeks
Now	38.4 weeks



#### Closing points

- Psychosocial indices differ by intention status with "Now" having a healthier psychosocial profile and "Never" having a less healthy psychosocial profile
  - Depression (CES-D)
  - Self-efficacy
  - Interpersonal Support
  - Paternal Support
  - Perceived Stress
  - Personality
- Pregnancy intention is a multifaceted construct.




# Acknowledgements

- National Institutes of Health
- U.S. Environmental Protection Agency
- Duke University and Medical Center
- Study participants
- Duke University Office of Research Support






# **Pregnancy Intention and Maternal Age**






# Race by Maternal Age




Non-Hispanic Blacks



Non-Hispanic Whites




# **Pregnancy Intention by Race**






# **Pregnancy Intention and Income**






# **Pregnancy Intention and Income**

#### Non-Hispanic Blacks versus Non-Hispanic Whites





# **Pregnancy Intention and Insurance**





# Paternal Relationship

