## References

- American Physical Therapy Association (2014). *Guide to Physical Therapist Practice 3.0*.

  Retrieved October 19, 2015, from American Physical Therapy Association:

  <a href="http://guidetoptpractice.apta.org/content/1/SEC1.body">http://guidetoptpractice.apta.org/content/1/SEC1.body</a>
- Anandacoomarasamy, A., Caterson, I., Sambrook, P., Fransen, M., & March, L. (2008). The impact of obesity on the musculoskeletal system. *International Journal of Obesity*, 211-222. doi:10.1038/sj.ijo.0803715
- Bandura, A. (1977). Self-efficacy: Toward a unifying theory of behavioral change.

  \*Psychological Review, 84(2), 191-215.
- Centers for Disease Control and Prevention. (2015a). *Physical Activity and Health*. Retrieved October 19, 2015, from Centers for Disease Control and Prevention:

  <a href="http://www.cdc.gov/physicalactivity/basics/pa-health/index.htm">http://www.cdc.gov/physicalactivity/basics/pa-health/index.htm</a>
- Centers for Disease Control and Prevention. (2015b). *Arthritis*. Retrieved October 19, 2015, from Centers for Disease Control and Prevention:

  <a href="http://www.cdc.gov/chronicdisease/resources/publications/aag/arthritis.htm">http://www.cdc.gov/chronicdisease/resources/publications/aag/arthritis.htm</a>
- Centers for Disease Control and Prevention. (2013). *Overweight and obesity for professionals:*Health consequences. Retrieved from Centers for Disease Control and Prevention:

  http://www.cdc.gov/obesity/adult/causes/index.html
- Chen, C., Neufeld, P., Feely, C., Skinner, C. (1999). Factors influencing compliance with home exercise programs among patients with upper-extremity impairment. *The American Journal of Occupational Therapy*, 53, 171-180.

- Finkelstein, E. A., Trogdon, J. G., Cohen, J. W., & Dietz, W. (2009). Annual medical spending attributable to obesity: Payer-and service-specific estimates. *Health Affairs*, 28(5), w822-w831. doi:10.1377/hlthaff.28.5w822
- Gyurcsik, N., Brawley, L., Spink, K., Brittain, D., Fuller, D., Chad, K. (2009). Physical activity in women with arthritis: Examining perceived barriers and self-regulatory efficacy to cope. *Arthritis and Rheumatism*, 61(8), 1087-1094.
- Josyula, L., Lyle, R. (2013). Health care provider physical activity prescription intervention. *American Journal of Health Education*, 44, 162-168.
- Kerssens, J., Sluijs, E., Verhaak, P., Knibbe, H., Hermans, I. (1999). Educating patient educators: enhancing instructional effectiveness in physical therapy for low back pain. *Patient Education and Counseling*, *37*, 165-176.
- Smeets, R., Vlaeyen, J., Hidding, A., Kester, A., van der Heijden, G., van Geel, A., Knottnerus J. (2006). Active rehabilitation for chronic low back pain: Cognitive-behavioral, physical, or both? First direct post-treatment results from a randomized controlled trial. 

  BMC Musculoskeletal Disorders, 7(5). doi:10.1186/1471-2474-7-5
- Turner, J., Holtzman, S., Mancl, L. (2007). Mediators, moderators, and predictors of therapeutic change in cognitive-behavioral therapy for chronic pain. *Pain*, 127, 276-286.
- United States Department of Health and Human Services. (2014, December 11). *Chapter 2:*\*Physical Activity Has Many Health Benefits. Retrieved from Office of Disease

  \*Prevention and Health Promotion:

  http://www.health.gov/paguidelines/guidelines/chapter2.aspx